



DATED: August 18, 2003
SCARBOROUGH STATION
SCARBOROUGH, NEW YORK 10510-0827
(914) 941-5600

RECEIVED

OCT 1 4 2003

TECHNOLOGY CENTER R3700

I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH
THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL IN AN ENVELOPE
ADDRESSED TO: COMMISSIONER OF PATENTS AND TRADEMARKS,
WASHINGTON, D.C. 20231, NO.: EV323630555US.

McGLEW AND TUTTLE, P.C., SCARBOROUGH STATION,
SCARBOROUGH, NEW YORK 10510-0827

BY:

Vernon B. Forte

DATE: August 18, 2003

RECEIVED
AUG 21 2003
TC 3100 MAIL ROOM

31. A method of improving bioavailability of ergot derivatives administered using sustained-release delivery systems comprising combining an ergot derivative or mixture thereof with a pharmaceutically acceptable hydrophilic swelling agent or mixture thereof and one or more pharmaceutically acceptable excipients, and wherein the bioavailability is at least 25% higher than the bioavailability of the ergot derivative or mixture thereof administered using a conventional drug delivery system;

said ergot derivative is selected from the group consisting of α -dihydroergocryptine and bromocriptine.

32. A sustained-release pharmaceutical composition comprising:
an ergot derivative or mixture thereof;
a pharmaceutically acceptable swelling agent or mixture thereof; and
one or more pharmaceutically acceptable excipients;

at cont,
said composition having a bioavailability at least equal to the bioavailability of the ergot derivative or mixture thereof administered using a conventional drug delivery system, and wherein said ergot derivative is selected from the group consisting of α -dihydroergocryptine and bromocriptine.

33. The composition according to Claim 32, wherein the bioavailability is at least 25% higher than the bioavailability of the ergot derivative or mixture thereof administered using a conventional drug delivery system.

34. The composition according to Claim 32, wherein the ergot derivative is α -dihydroergocryptine.

35. The composition according to Claim 32, wherein the hydrophilic swelling agent is selected from the group consisting of methylcellulose, carboxymethylcellulose, hydroxypropylmethylcellulose, polyvinyl alcohols, polyoxyethylene glycols and poloxamers and mixtures thereof.